
Role of 11beta-hydroxysteroid Dehydrogenase Inhibitors in Metabolic Syndrome and its Expansion in other Therapeutic Options

Raj Kapoor B,* Ali Dau Masoud,* Abdussalam Sughir,** and Vijayakumar A,***

Abstract

The metabolic syndrome is a constellation of interrelated metabolic risk factors that appear to promote the development of diabetes and cardiovascular disease. These risk factors include abdominal obesity, insulin resistance, hypertension and dyslipidemia. 11beta-Hydroxysteroid dehydrogenase (11beta-HSD) catalyzes the interconversion of glucocorticoids through the activity of two isozymes: type 1 (11beta-HSD1) and type 2 (11beta-HSD2). 11beta-HSD1 converts inactive glucocorticoid to the active form, whereas 11beta-HSD2 converts active glucocorticoid to the inactive form. Glucocorticoids play a pivotal role in regulating fat metabolism, function and distribution. Evidence has accumulated that enzyme activity of 11beta-hydroxysteroid dehydrogenase type 1 (11beta-HSD1), which regenerates active glucocorticoids from inactive forms and plays a central role in regulating intracellular glucocorticoid concentration, is commonly elevated in fat depots from obese individuals. This suggests a role for local glucocorticoid reactivation in obesity and the Metabolic Syndrome. 11beta-HSD1 knockout mice resist visceral fat accumulation and insulin resistance even on a high-fat diet. In summary, 11beta-HSD1 is a promising pharmaceutical target for the treatment of the Metabolic Syndrome. Animal studies and pharmacological experiments suggest further unrelated target areas, for example improvement of cognitive function and treatment of glaucoma, osteoporosis due to the role of glucocorticoids and cellular activation by 11 beta HSD1 in these pathologies.

KEY WORDS: 11 Beta hydroxysteroid dehydrogenase inhibitors, metabolic syndrome, glucocorticoids, obesity, type 2 diabetes.

1. Introduction

Glucocorticoids (GCs) are steroid hormones secreted by the adrenal glands in a circadian fashion, further modulated by ectopic stress signals such as trauma, inflammation, psychological stress or physical exercise.¹⁻³

These hormones affect every tissue and organ in the body and thus contribute to maintain overall homeostasis and, importantly, are one of the most potent modulators of the stress response in mammals. Besides the stress mediated effects of the sympathetic nervous system on metabolic alterations that contribute to the etiology of diabetes mellitus, atherosclerosis and cardiovascular disease,⁴ GCs are among the critical components identified in metabolic disease.

Secretion of GCs from the adrenal cortex is controlled by negative feedback via the hypothalamic pituitary adrenal (HPA) axis. The main regulators of intracellular GC levels are 11 beta hydroxysteroid dehydrogenase (HSD) enzymes. Two isoforms of 11 beta HSD have been cloned and characterized.^{5,6}

11beta HSD type 1 is an NADP(H)-dependent enzyme that acts primarily as a reductase in intact cells, converting the inactive 11-keto metabolites cortisone (in humans) or 11-dehydrocorticosterone (in rodents) into the active GCs cortisol or corticosterone, respectively. 11 beta HSD1 is expressed in most tissue types and potentiates the action of endogenous GCs by increasing their local concentration. 11 beta HSD type 2 is an NAD (H)-dependent enzyme that catalyzes the reverse reaction, oxidizing active GCs to their inactive 11-keto forms. Although 11 beta HSD1 is widely expressed, 11 beta HSD2 expression is limited to tissues that express the mineralocorticoid receptor, such as the kidney and gut, as well as to the placenta. By inactivating cortisol, 11 beta HSD2 prevents it from binding to the mineralocorticoid receptor, thus conferring aldosterone specificity on the receptor. In the placenta, the enzyme prevents maternal GCs from reaching the fetal circulation.

*) Department of Pharmacology, Faculty of Medicine, Sebha University, Sebha, Libya

**) Faculty of Pharmacy, University of Almergib, Alkhoms, Libya.

***) Drug and Poison Information Center, Department of Pharmacy Practice, KMCH College of Pharmacy, Kalapatti Road, Coimbatore 641 048, India.

The metabolic syndrome is a cluster of cardiovascular risk factors, including visceral obesity, insulin resistance, dyslipidemia, and hypertension. It has been suggested that metabolic syndrome may result from increased intracellular GC tone, it may occur with elevated 11beta -HSD1 activity, and pharmacologic inhibition of 11 beta -HSD1 may alter intracellular GC levels and might be therapeutic option for metabolic syndrome.⁷⁻⁹

2. Role of Glucocorticoids in Metabolic Syndrome

The critical role of GCs for mammalian homeostasis is clearly highlighted by the phenotypical changes observed in deficiency to produce these steroids (Addison's disease) and those observed in situations of overproduction (Cushing's disease). Adrenal insufficiency to produce GCs (caused by destruction of the adrenal gland e.g. by infection, autoimmune disorders, tumor metastases) leads to general weakness, pigmentation of the skin, weight loss and hypotension. In contrast, the systemic high levels of GCs observed in patients with Cushing's syndrome (usually with a pituitary tumor leading to overproduction of ACTH, the hormone controlling GC release and production from the adrenal gland or through pharmacological immunosuppressive treatment with synthetic GCs as in inflammatory disorders or in transplantation settings) show central obesity, insulin resistance, glucose intolerance, behavioral changes, dyslipidemia, and hypertension. The main metabolic effects of GCs in respect to the metabolic syndrome comprise (as the name implies) regulation of fasting blood glucose levels (through hepatic induction of gluconeogenesis enzymes, mainly glucose-6-phosphatase and phosphoenolpyruvate carboxykinase (PEPCK), control of adipocyte proliferation, differentiation and metabolism, control of skeletal muscle insulin sensitivity, and regulation of insulin secretion from the beta-cells of the endocrine pancreas.

The metabolic changes occurring in Cushing's disease are also observed in patients with the metabolic syndrome and constitute independent risk factors for the development of cardiovascular disease,¹⁰⁻¹³ still among the most health threatening conditions in western societies and rapidly progressing in third world countries. Importantly, these symptoms are usually reversible in Cushing's disease, upon correction of the primary cause. However, despite extensive efforts, no systemic elevation of GC levels is observed in patients with the metabolic syndrome, an enigma that is solved

by the discovery that the GC activating enzyme 11beta HSD1 is present in adipose tissue or liver, an effect, which locally produces elevated tissue levels of GCs, in the light of normal systemic GC levels.^{14,15}

3. 11beta-Hydroxysteroid Dehydrogenase Type 1 as an Amplifier of Glucocorticoid Action

More recently, the focus of attention has moved to 11beta-hydroxysteroid dehydrogenase type 1 (11HSD1). Although this enzyme can catalyze dehydrogenase conversion from cortisol to cortisone *in vitro* when deprived of regeneration of reduced nicotinamide adenine dinucleotide phosphate (NADPH) cofactor, *in vivo* it functions as a predominant, if not exclusive, reductase converting cortisone to cortisol.¹⁶ 11HSD1 is most highly expressed in liver, adipose tissue, lung, and areas of CNS.¹⁵ In these sites it is associated with glucocorticoid, not mineralocorticoid, receptors. Because glucocorticoid receptors have relatively low affinity for cortisol, and cortisone levels in blood are relatively constant and even exceed the free levels of cortisol during the diurnal nadir, we hypothesized that the physiological role of 11HSD1 is to maintain adequate activation of glucocorticoid receptors in sites where this is vital to metabolic function.¹⁷

It follows that increased 11HSD1 activity may generate increased cortisol levels within adipose tissue and liver and thereby promote features of the metabolic syndrome, and that inhibition of 11HSD1 may lower cortisol levels within these tissues and hence be beneficial in the metabolic syndrome. Such changes in tissue cortisol concentrations could occur without any changes in plasma cortisol levels provided the latter are maintained by normal feedback regulation of the HPA axis.

4. 11beta-hsd1 as a potential drug target in the metabolic syndrome

One of the most important discoveries in 11beta -HSD1 research was that the enzyme functions *in vivo* or in cell culture primarily as reductase thus amplifying intracellular cortisol levels.¹⁸⁻²¹ This was further elegantly corroborated by the 11beta-HSD1 knockout mouse (11beta-HSD1-/-),²² showing that the enzyme is the sole 11-ketosteroid reductase, at least in mice. A logical translation of this concept into first clinical trials using the 11beta-HSD isozyme-unselective inhibitor carbenoxolone showed that 11beta-HSD inhibition indeed might lead to increased hepatic insulin sensitivity in healthy

volunteers,^{23,24} but also in a small set of diabetic patients. These initial analyses delivered for the first time substantial evidence that inhibition of 11beta HSD1 might be a useful strategy in the treatment of metabolic disorders such as insulin resistance and possibly obesity. The 11beta-HSD1^{-/-} animals are fertile, reproduce normally and exhibit a normal lifespan.²² A diabetes-resistant phenotype is only apparent upon challenging by a high-fat diet or stress, showing attenuated responses to these stimuli, with altered glucose levels, fat accumulation and lipid profiles.^{22,25,26} The hypothalamic-pituitary-adrenal (HPA) axis appears to be disturbed as shown by adrenal hyperplasia, indicating that cortisone reduction by 11beta-HSD1 is also important in the CNS feedback control and HPA axis regulation.^{22,27} Later on, transgenic mice were created with a rat 11beta-HSD1 gene under the control of the promoter of the adipocyte fatty acid binding protein 2 (aP2-11beta-HSD1 mouse);²⁸ resulting in 2-3 fold elevated levels of 11beta-HSD1 in visceral adipocytes. This expression level is also observed in several²⁹⁻³³ but not all studies of obese patients³⁴⁻³⁷ or in genetic rodent models of obesity such as the Zucker rat.^{22,27} Association studies suggest a correlation between GC metabolism and obesity markers such as body mass index or waist to hip ratio.³⁸⁻⁴⁰

A further indication for the significance of 11beta-HSD1 and GCs in visceral fat accumulation is a Cushing patient case with defective peripheral cortisone to cortisol conversion, displaying a non obese phenotype.⁴¹ The aP2-11beta-HSD1 mice have an about 2-fold concentration level of adipose tissue corticosterone as compared to wild-type animals, whereas plasma GC levels are normal. Importantly, these aP2-11beta HSD1 animals show the cardinal symptoms of the metabolic syndrome. They are obese, predominantly through visceral fat accumulation achieved through differential GR distribution in visceral and subcutaneous adipocytes, and display insulin resistance, glucose intolerance, leptin resistance, have elevated free fatty acid and triglyceride levels, and are hypertensive.^{29,42} The arterial hypertension is an effect obtained through activation of the renin-angiotensin-aldosterone system, likely driven by elevated levels of adipocyte derived angiotensinogen.^{29,42} Altered expression levels compared to controls for key molecules such as leptin, adiponectin,

resistin, TNF-alpha, UCP and PPAR-gamma were observed.^{29,42} A non-obese phenotype of the metabolic syndrome but with accompanying dyslipidemia and arterial hypertension, caused by hepatic activation of the renin-angiotensin system, was achieved through hepatic overexpression of 11 beta HSD1 using the apoE promoter (apoE-11beta-HSD1 mouse).⁴³ These animals displayed fatty liver and increased hepatic fatty acid synthesis and fatty acid beta-oxidation by upregulation of LXR-alpha and PPAR-alpha mRNA levels, resulting in impaired lipid clearance.⁴³ This animal line might be useful as a model system for myotonic dystrophia or metabolically obese, but normal weight individuals.^{43,44} Hepatic inhibition of 11beta HSD1 achieved in several mouse models of insulin resistance by a first series of selective 11beta HSD1 inhibitors developed by Biovitrum in Stockholm.⁴⁵⁻⁴⁷ Down regulation of hepatic gluconeogenic enzymes like PEPCK and glucose 6 phosphatase was accompanied by a significant decrease of insulin and glucose levels, as well as lowered cholesterol, triglyceride and free fatty acid levels.⁴⁵⁻⁴⁷ Accordingly, these results demonstrated that enzyme inhibition in the liver leads to improved insulin sensitivity and lipid status in the ob/ob, db/db and KKAY animal model systems used.^{46,47}

5. Expression of 11 Beta Hsd1 in Skeletal Muscle

The expression of 11beta HSD1 was not only in liver and adipose tissue, but also it was in skeletal muscle.⁴⁸⁻⁵⁰ The endocrine pancreas⁵¹ suggests a critical role of this GC activating system for regulation of overall insulin sensitivity. In skeletal muscle, the interplay between GR alpha and beta isoforms and the expression of 11beta-HSD1 in a study using human myoblast cultures, and the potential consequences for overall insulin sensitivity were presented.^{48,49} Interestingly insulin was appeared to upregulate 11beta-HSD1 expression level in the presence of unidentified serum factors.^{48,49} thereby constituting a feedback control system, in contrast to the insulin effects observed on 11beta-HSD1 expression levels in liver.^{20,21} Expression of 11beta-HSD1 in skeletal myoblasts was associated with insulin, increased body mass index (BMI) and blood pressure.^{48,49} Taken together, the studies supported the notion that skeletal muscle 11beta-HSD1 contributes to features of the metabolic syndrome and that pharmacological inhibition of 11beta-HSD1

might lead to increased insulin sensitivity mediated in part by increased triglyceride uptake in skeletal muscle.⁵⁰

6. 11 Beta Hsd1 in Pancreatic Cells

The presence of 11beta- HSD in pancreatic islets and beta-cells was first established in the ob/ob mouse system.⁵¹ The fundamental role of GC on beta-cell physiology was investigated in several studies, showing that increased glucocorticoid sensitivity leads to inhibition of insulin release.^{52,53} Transgenic mice, overexpressing the GR in beta-cells under the control of the insulin promoter show increased GC sensitivity, and animals ultimately develop diabetes with a decreased capacity in insulin secretion.^{52,53} In several cell culture systems, GCs display profound and complex effects on insulin secretion, with enhancing or inhibiting effects, depending on the experimental conditions.^{51,52,55-57} In ob/ob mice GCs inhibit insulin secretion,^{51,58} further amplified by 11 beta HSD1.⁵¹ Blockade by CBX or synthetic selective 11beta-HSD1 inhibitors such as the aryl sulfonamide thiazole BVT2733 enhances glucose stimulated insulin secretion (GSIS).^{51,58} Therefore 11beta-HSD1 might constitute an essential factor for the ability of the endocrine pancreas to adequately secrete insulin, and accordingly could play an effective role in the pathogenesis of insulin resistance.

A convincing proof of concept that 11beta-HSD1 is a valid target for treatment of the metabolic syndrome was derived from animal models treated with a selective 11beta-HSD1 adamantyltriazol inhibitor developed by Merck.^{12,59,60} The compound potently inhibits 11beta-HSD1 activity in liver and importantly also in adipose tissue. In diet induced obesity, the compound lowers weight gain, food intake and fat accumulation, and similar to the effects observed with other inhibitors improves insulin sensitivity and lipid profile, e.g. cholesterol, triglyceride and free fatty acid levels.¹² Furthermore, in the apoE knockout mouse, the compound dramatically reduces atherosclerotic plaque formation, and for the first time suggests direct beneficial effects of enzyme disease.¹² Since the lipid effects observed were mainly a moderate decrease in lipid levels, these actions on plaque formation are likely to be mediated through direct effects on the complex arteriosclerotic process.^{12,61} Importantly, 11betaHSD1 expression has been demonstrated in endothelial cells, activated macrophages and smooth muscle cells, and likely involves a yet to be defined role of

11beta HSD1 in inflammatory mediation of plaque formation.^{12,61} Taken together, all animal and pharmacological studies carried out thus far support the concept that blockade of 11beta HSD1 might be a valid, novel approach in the treatment of all facets of the metabolic syndrome.

7. Potential Effects of 11HSD1 Inhibitors beyond Adipose Tissue and Liver

The widespread distribution of 11HSD1 raises the possibility that target related toxicity will occur in nonmetabolic tissues. For example, 11HSD1 is increased in differentiation of monocytes to macrophages,⁶² and lack of glucocorticoid action within macrophages may be deleterious in inflammation. 11HSD1 is also expressed in vascular smooth muscle.⁶³ Although 11HSD1 null mice have normal vascular contractile and relaxant function⁶⁴ recently finding shown that lack of glucocorticoid regeneration within the vessel wall removes an angiostatic "brake."⁶⁵ As a result, enhanced angiogenesis may occur during 11HSD1 inhibition which may be beneficial in healing of wounds, and may even improve myocardial recovery after ischemia,⁶⁵ but might also allow unrestrained angiogenesis, for example, in diabetic retinopathy. The expression of 11HSD1 in the central nervous system, particularly in hippocampus, offers both opportunities and threats for the success of 11HSD1 inhibitors.

The concern has been that inhibition of hippocampal 11HSD1 may reduce negative feedback suppression of the HPA axis, resulting in a compensatory increase in plasma cortisol which overcomes the benefits of peripheral 11HSD1 inhibition. However, although 11HSD1 null mice do have a subtle elevation in plasma glucocorticoid levels,⁶⁶ they retain the beneficial effects of 11HSD1 inhibition in peripheral tissues. 11HSD1 null mice are protected from cognitive dysfunction with aging⁶⁷ and administration of carbenoxolone to humans enhances cognitive performance⁶⁸ on balance, inhibition of 11HSD1 in CNS appears to be beneficial rather than deleterious.

8. Role of 11HSD1 in Inflammatory Regulation

Inflammatory cytokines, such as interleukin-1 (IL-1) and tumor necrosis factor - α (TNF- α), upregulate 11HSD1 expression in a number of different cells [14]. In monocytes, 11HSD1 increases in association with differentiation into competent phagocytic macrophages.²⁵

These observations are consistent with 11HSD1 playing a role in amplifying local glucocorticoid concentrations within the site of injury or infection, operating in addition to the activation of the HPA axis which accompanies systemically important inflammation.⁶⁹⁻⁷² Obesity has been described as an inflammatory state, and it is possible that increased activity of immune cells within adipose tissue underlies the upregulation of 11HSD1 which accompanies obesity,⁷³⁻⁷⁵ This remains, however, a speculation.

9. Nutritional Regulation of 11HSD1

A number of hormones which respond to nutritional status also influence 11HSD1, including insulin,^{27,76} growth hormone and insulin-like growth factor-1 (IGF1).²⁷ In addition, peroxisome-proliferator activated receptor (PPAR)^{29,30} and liver X receptor (LXR) agonists³¹ alter 11HSD1 expression, consistent with a response to variations in fatty acids and/or oxysterols. Furthermore, variations in glucose flux may affect H6PDH activity and hence 11HSD1 equilibrium. Few direct experiments to test the effects of nutritional status on 11HSD1 *in vivo* have been published. Morton *et al.* showed that high fat feeding of mice down regulated 11HSD1 in adipose tissue.³¹ Leptin administration to ob/ob mice also upregulates liver 11HSD1, although it is not clear whether this is a direct effect of leptin or an indirect effect of weight loss.⁷⁷ In humans, weight loss has variably been reported to increase⁷⁸ or decrease⁷⁹ adipose 11HSD1; it is possible that this discrepancy reflects the different dietary exposure of subjects at the time that biopsies were taken. In mice, sensitivity of 11HSD1 to regulation by dietary fat predicts susceptibility to diet-induced obesity.³² It is an intriguing possibility that the primary purpose of 11HSD1 is to facilitate appropriate adaptations to variations in diet, and that dysregulation of this response underlies increased 11HSD1 in adipose tissue in obesity.

10. Further indications for selective 11 beta HSD1 inhibition

Besides the potential beneficial use of selective 11 beta HSD1 inhibitors to treat insulin resistance, obesity, dyslipidemia and cardiovascular disease, several studies suggest further potential applications, as outlined in the following subchapters. Some of these applications, namely the GC amplification effects by 11 beta HSD1 on the CNS and in glaucoma, appear to be rather validated, whereas several others, e.g. the role of 11 beta

HSD1 in immunity and inflammatory processes clearly deserve further investigation and experimental efforts.

11. Glucocorticoids and the CNS: the Role of 11 beta HSD1 in Ageing and Neurodegeneration:

GCs exert a multitude of effects on the central nervous system affecting among others neurotransmitters, receptors, ion channels and metabolism.⁸⁰⁻⁸² Chronically elevated levels of GCs clearly impair neuropsychiatric functions and can lead to dysfunctions such as depression, anxiety, psychosis or euphoria. Thus considerable evidence exists which links GCs with cognitive functions and the neuroendocrine axis.⁸⁰⁻⁸² A highly sensitive site for GC effects is the hippocampus, a brain structure necessary to integrate multiple neuroendocrine, cognitive and affective signals. Expression of 11 beta HSD1 in the central nervous system was shown to be mainly restricted to the hippocampus, cerebellum and the cortex.⁸³⁻⁸⁵ The first experiments demonstrating a possible role of 11 beta HSD1 inhibition in blocking neurotoxicity were performed using primary cultured cells derived from fetal rat hippocampus, and it was shown that CBX inhibition of 11 beta HSD1 was able to prevent kainic acid induced neurotoxicity.⁸⁶ Further insight was obtained from 11 beta HSD-/- animals, showing that aged wild-type animals show GC associated, hippocampal driven, significant learning and memory impairments in the water maze test.⁸⁷ Importantly, aged knock-out animals perform not significantly different in these tests than young wild-type or young knockout animals, suggesting that hippocampal 11 beta HSD1 inhibition might show significant effects and protects against ageing related loss of cognitive functions.⁸⁷ Indeed, in a double-blind, placebo controlled crossover study with healthy volunteers and diabetic patients (age matched, 52-75 years of age), administration of CBX (100 mg/ 3 times/day) resulted in significant improvement of verbal fluency and memory function after 4 weeks of treatment[88]. The recently described rare haplotype discovered in the regulatory region of the HSD11B1 gene⁸⁹ was associated with a 6-fold increased risk for sporadic Alzheimer's disease, adding a further piece of evidence to the role of GCs and their metabolic activation in cognitive function. Importantly, an undisclosed compound from a recently developed selective inhibitor series by

Merck (unpublished data) shows similar positive effects in animals subjected to passive avoidance or novel object recognition tests, suggesting that applications of selective 11 beta HSD1 inhibitors indeed go well beyond the treatment of the metabolic syndrome.

12. Role of 11 beta HSD1 in Glaucoma

Increase in intraocular pressure can be provoked by ocular or systemic administration of GCs, and furthermore, patients with primary open angle glaucoma exhibit increased GC sensitivity.^{90,91} Expression of 11 beta HSD1 has been shown in nonpigmented epithelial cells of the ciliary body, the site of aqueous humor production,^{90,91} likely to be the reason for the observed increased cortisol:cortisone ratio in humor. Topical administration of the 11 beta HSD inhibitor CBX lowers intraocular pressure in healthy volunteers and patients with ocular hypertension, pointing to a potential novel application for 11 beta HSD1 inhibitors.^{90,91}

13. Inflammation and Immunity: What is the Function of 11 beta HSD1 in the Immune System?

Since their introduction into clinical medicine GCs almost 50 years ago⁹² GCs have been a cornerstone in the treatment of chronic inflammatory diseases such as rheumatoid arthritis, asthma bronchiale and autoimmune diseases are used in transplantation settings to prevent graft versus host disease.^{1-3,94} Consequently, much has been learned about the underlying pathophysiological mechanisms by the use of synthetic GCs. GCs exert potent effects on the immune system and it now appears to be oversimplified that GCs have merely an anti-inflammatory or immunosuppressive profile.^{95,96} The dogmatic viewpoint of immunosuppressive GC action appears to be supported by the fact that GCs have been mainly used in pharmacological i.e. supraphysiological doses. Many recent studies however suggest that the role of GCs on the immune system is extremely complex and shows a time, dose, compound and context dependent pattern of reactions on critical cells of the immune system.^{95,96} It appears that inflammatory processes are regulated in a time dependent manner and controlled by synergistic and antagonistic actions of GCs and cytokines. As discussed in two overviews^{95,96} GCs upregulate the expression of IL-6 receptors, the signal transduction component gp13, of IL-1 receptors or can promote release of inflammatory signals such as TNF α , IL-6 or macrophage migration

inhibitory factor.^{95,97} Thus it appears that GC effects are rather immunoregulatory than only immunosuppressive. Furthermore, GCs are intimately linked to their capacity to alter specifically the secretion pattern of cytokines associated with inflammatory and immune processes.⁹⁵ GCs modify the Th1-Th2 cytokine profile by favouring a Th2 profile and might represent an important determinant for the development of a specific Th1/Th2 pattern in inflammatory situations. Not unexpectedly, 11 betaHSD1 is expressed at important sites of inflammatory responses. Whereas peripheral lymphocytes appear to be devoid of 11 beta HSD isozymes,⁹⁸ enzymatic activity and expression of 11 beta HSD1 was noted in stromal cells of lymphoid tissues[99] , in activated monocytes or in dendritic cells.¹⁰⁰ Thus far the use of inhibitors of

11 beta HSD enhanced the local effects of GCs in patients with psoriasis or eczema¹⁰¹ or decreased resistance to experimental *Listeria* infections.¹⁰² Taken together, modulation of local activation of GC activation in immune tissues by 11 beta HSD inhibitors might offer the exciting possibility to selectively drive and direct immune responses,^{95,96,100} but final proof of concept for this hypothesis is still missing and experiments in several laboratories are currently underway.

14. Glucocorticoid Induced Osteoporosis: Does 11 beta HSD1 Play a Role?

Chronic exposure of endogenous and synthetic GCs has fundamental effects on bone structure and function,¹⁰³ with osteopenia, osteoporosis and increased risk for bonefractures as well recognized consequences. These detrimental effects are potentially reversible at least in part, and amenable to treatment e.g. by use of bisphosphonates.¹⁰⁴ GCs exert multiple effects on bone cells, by increasing bone resorption and decreasing bone formation, however the underlying mechanisms are at present far away from being completely understood. The potential impact of 11 beta HSD expression in bone cells^{13,105-107} was addressed in some studies. Short-term inhibition with CBX in healthy volunteers had no effect on bone formation markers, whereas a suppression of bone resorption markers was noted.¹⁰⁶ It was proposed that expression of 11 beta HSD1 in osteoblasts affects proliferation and differentiation and due to predominant reductase activity in intact cells constitutes a risk factor in age related decrease in bone formation and increased risk for GC induced osteoporosis.¹⁰⁶ Somewhat surprising was the

lack of a clear bone phenotype in the 11 beta HSD1-/- mouse model. The animals exhibited no significant changes in cortical or trabecular bone mass, and aged animals show similar levels of bone loss compared to matched control animals,¹⁰⁷ however, a complete absence of bone marrow adipocytes was noted. Clearly further studies are necessary to determine the role of 11 beta HSD isozymes and potential pharmacological intervention of 11 beta HSD inhibition in bone tissue

15. Perspectives and Commentary

As discussed in this overview selective inhibition of 11beta HSD1 could in principle offer unprecedented and unique opportunities to improve all symptoms of the metabolic syndrome, which taken together, constitute independent risk factors for the development of cardiovascular disease. Due to the central role of GCs in control of metabolism and the function of 11beta-HSD1 as a pre-receptor control of the GR, tissue selective blockade could be a general principle to improve insulin sensitivity, lower lipid levels, regulate fat accumulation and decrease arterial hypertension, the cardinal features of the metabolic syndrome. Clearly, further innovative approaches are needed besides the established pharmacological principles of e.g. PPAR alpha and PPAR gamma agonists to lower lipids or increase insulin sensitivity in order to combat the consequences of the world wide dramatic increase in obesity and diabetes. In fact, several of the beneficial effects observed with fibrates or thiazolidine-diones might be due to their potential to down regulate 11beta-HSD1 in liver or adipose

tissue. Moreover, the tight connection between lipid and glucose metabolism^{72,73} might be linked through the LXR-GR-11betaHSD1 axis.⁷¹ Blockade of the hepatic GR to increase insulin sensitivity and lower blood glucose have been shown to be a valid concept in the treatment of diabetes⁷⁴⁻⁷⁷ and at least in several animal models a variation of this theme by inhibition of hepatic 11beta-HSD1 has been shown effective.^{46,47} Targeting of hepatic 11beta-HSD1 might thus be a valid principle in lean diabetic individuals or in a subset of obese patients with elevated hepatic GC reamplification.²⁶ However, in several, but not all studies of human obesity hepatic expression of 11beta-HSD1 is apparently down regulated and might therefore not be the primary target for treatment of insulin resistance and hyperglycemia.⁷⁷ Immunogenic 11beta-HSD1 models, adipocyte 11beta-HSD1 plays the central role in mediating features of the metabolic syndrome and therefore might constitute the prime pharmacological target. Human idiopathic metabolic syndrome is undoubtedly more complex in terms of GC amplification.^{29,34,35,37} This is further corroborated by a recent mouse model of obesity and insulin resistance,²⁶ with elevated hepatic 11beta-HSD1 activity but decreased adipose 11beta-HSD1 levels, indicating that such a pattern of GC amplification might exist in human idiopathic obesity as well. Hence it is clearly mandatory to investigate further patient populations to illuminate the role of GC amplification by 11beta-HSD1, determine the status of 11beta-HSD1 expression and address the suitability of 11beta-HSD1 inhibitor therapy.

References:

1. Adcock I M, Cosio B, Tsaprouni L, Barnes PJ, Ito K. Redox regulation of histone deacetylases and glucocorticoid-mediated inhibition of the inflammatory response. *Antioxid. Redox. Signal* 2005;7:144-52.
2. Adcock I M, Ito K, Barnes P J. Glucocorticoids: effects on gene transcription. *Proc. Am. Thorac. Soc* 2004; 1: 247-54
3. Barnes P.J, Adcock I. Anti-inflammatory actions of steroids: molecular mechanisms. *Trends Pharmacol. Sci.* 1993;14: 436-41
4. Nonogaki K, Iguchi A. Role of central neural mechanisms in the regulation of hepatic glucose metabolism. *Life Sci.* 1997; 60: 797-807.
5. Agarwal A K, Monder C, Eckstein B, White P C. Cloning and expression of rat cDNA encoding corticosteroid 11 beta-dehydrogenase. *J. Biol. Chem.* 1989; 264: 18939-43
6. Stewart P M. Tissue-specific Cushing's syndrome, 11beta-hydroxysteroid dehydrogenases and the redefinition of corticosteroid hormone action. *Eur. J. Endocrinol.* 2003; 149: 163-68.
7. Albiston A L, V R Obeyesekere, R E Smith, Z S Krozowski. Cloning and tissue distribution of the human 11 beta-hydroxysteroid dehydrogenase type 2 enzyme. *Mol. Cell. Endocrinol.* 1994;105:R11-17.

8. Walker B R, J.R Seckl. 11beta-hydroxysteroid dehydrogenase type 1 as a novel therapeutic target in metabolic and neurodegenerative disease. *Expert Opin. Ther. Targets.* 2003; 7:771–83
9. Thieringer R, Hermanowski-Vosatka A. Inhibition of 11beta-HSD1 as a novel treatment for the metabolic syndrome: do glucocorticoids play a role?. *Exp. Rev. Cardiovasc. Ther.* 2005; 3: 911-24.
10. Bahr V, Pfeiffer A.F, Diederich S. The metabolic syndrome X and peripheral cortisol synthesis. *Exp. Clin. Endocrinol. Diabetes.* 2002;110: 313-18
11. Masuzaki H, J.S. Flier. Tissue-specific glucocorticoid reactivating enzyme, 11 beta-hydroxysteroid dehydrogenase type 1 (11 beta-HSD1)—a promising drug target for the treatment of metabolic syndrome. *Curr. Drug Targets Immune Endocr. Metabol. Disord.* 2003;3:255–62.
12. Thieringer R, Hermanowski-Vosatka, A. Inhibition of 11beta-HSD1 as a novel treatment for the metabolic syndrome: do glucocorticoids play a role?. *Exp. Rev. Cardiovasc. Ther.* 2005; 3 :911-924.
13. Tomlinson J.W, Walker E.A, Bujalska I.J, Draper N, Lavery G.G, Cooper M.S, Hewison M, Stewart P.M. 11β-hydroxysteroid dehydrogenase type 1: A tissue-specific regulator of glucocorticoid response. *Endocr. Rev.* 2005; 25: 831-66
14. Ulick S, Levine LS, Gunczler P, et al. A syndrome of apparent mineralocorticoid excess associated with defects in the peripheral metabolism of cortisol. *J Clin Endocrinol Metab* 1979; 49:757-64.
15. Walker B R, Connacher A A, Lindsay R M, Webb D J, Edwards C R. Carbenoxolone increases hepatic insulin sensitivity in man: a novel role for 11-oxosteroid reductase in enhancing glucocorticoid receptor activation. *J. Clin. Endocrinol. Metab.* 1995; 80: 3155–59
16. Seckl JR, Walker BR. 11beta Hydroxysteroid dehydrogenase type 1—a tissue-specific amplifier of glucocorticoid action. *Endocrinology.* 2001;142: 1371–76
17. Jamieson PM, Walker BR, Chapman KE, Andrew R, Rossiter S, Seckl JR. 11beta-hydroxysteroid dehydrogenase type 1 is a predominant 11 beta-reductase in the intact perfused rat liver. *J Endocrinol.* 2000;165: 685-92
18. Low S C, Chapman KE, Edwards C R, Seckl J R. 'Liver-type' 11 beta-hydroxysteroid dehydrogenase cDNA encodes reductase but not dehydrogenase activity in intact mammalian COS-7 cells. *J. Mol. Endocrinol.* 1994;13:167-74.
19. Jamieson P M, Chapman K E, Edwards C R, Seckl J R. 11β-Hydroxysteroid dehydrogenase is an exclusive 11β-reductase in primary cultured rat hepatocytes: effect of physicochemical and hormonal manipulations. *Endocrinology.* 1995;136: 4754-61
20. Voice M W, Seckl J R, Edwards C R, Chapman K E. 11β-hydroxysteroid dehydrogenase type 1 expression in 2S FAZA hepatoma cells is hormonally regulated: a model system for the study of hepatic glucocorticoid metabolism *Biochem. J.* 1996; 317:621-25
21. Kotelevtsev Y, Holmes M C, Burchell A, Houston P M, Schmöll D , Jamieson P, et al 11β-hydroxysteroid dehydrogenase type 1 knockout mice show attenuated glucocorticoid-inducible responses and resist hyperglycemia on obesity or stress. *Proc. Natl. Acad. Sci. USA.* 1997; 94:14924-129.
22. Hewitt KN, EA Walker, P.M. Stewart. Hexose-6-phosphate dehydrogenase and redox control of 11 beta hydroxysteroid dehydrogenase type 1 activity. *Endocrinology.* 2005;146: 2539–43
23. Banhegyi G, A Benedetti, R Fulceri, S. Senesi. Cooperativity between 11beta-hydroxysteroid dehydrogenase type 1 and hexose-6-phosphate dehydrogenase in the lumen of the endoplasmic reticulum. *J. Biol. Chem.* 2004;279: 27017–21
24. Atanasov AG, LG Nashev, RAS Schweizer, et al. Hexose-6-phosphate dehydrogenase determines the reaction direction of 11beta-hydroxysteroid Dehydrogenase type 1 as an oxoreductase. *FEBS Lett.* 2004;571: 129–33.
25. Thieringer R, Le Grand CB, Carbin L, Cai TQ, Wong B, Wright SD, Hermanowski-Vosatka A. 11beta-Hydroxysteroid dehydrogenase type 1 is induced in human monocytes upon differentiation to macrophages. *J. Immunol.* 2001;167: 30–35
26. Hammani MM, PK Siiteri. Regulation of 11 beta hydroxysteroid dehydrogenase activity in human skin fibroblasts: enzymatic modulation of glucocorticoid action. *J. Clin. Endocrinol. Metab.* 1991;73: 326–34
27. Moore JS, JP Monson, G Kaltsas, et al. Modulation of 11 beta hydroxysteroid dehydrogenase isozymes by growth hormone and insulin-like growth factor: in vivo and in vitro studies. *J. Clin. Endocrinol. Metab.* 1999;84: 4172– 77

28. Trainer PJ, Drake WM, Perry LA, Taylor NF, Besser GM, Monson JP. Modulation of cortisol metabolism by the growth hormone receptor antagonist pegvisomant in patients with acromegaly. *J. Clin. Endocrinol. Metab.*2001; 86: 2989–92
29. Berger J, Tanen M, Elbrecht A, Hermanowski-Vosatka A, Moller D E, Wright S D. Peroxisome proliferator activated receptor- ligands inhibit adipocyte 11beta-hydroxysteroid dehydrogenase type 1 expression and activity. *J. Biol. Chem.*2001; 276: 12629–35
30. Hermanowski-Vosatka A, D Gerhold, SS.Mundt, et al.. PPARalpha agonists reduce 11beta-hydroxysteroid dehydrogenase type 1 in the liver.*Biochem.Biophys. Res. Commun.* 2000; 279: 330–36.
31. Stulnig, T M, U Oppermann K R, Steffensen, et al. PPARalpha agonists reduce 11beta-hydroxysteroid dehydrogenase type 1 in the liver. *Biochem. Biophys. Res. Commun.* 2002; 279: 330–36.
32. Morton NM, L Ramage, JR Seckyl. Down-regulation of adipose 11 beta hydroxysteroid dehydrogenase type 1 by high-fat feeding in mice: a potential adaptive mechanism counteracting metabolic disease. *Endocrinology.* 2004; 145: 2707–12.
33. Bujalska IJ, Walker E.A, Hewison M, Stewart PM. A switch in dehydrogenase to reductase activity of 11-β hydroxysteroid dehydrogenase type 1 upon differentiation of human omental adipose stromal cells. *J.Clin.Endocrinol. Metab.*2002;87:1205-10.
34. Bujalska IJ, Walker EA, Tomlinson JW, Hewison M, Stewart PM. 11Beta-hydroxysteroid dehydrogenase type 1 in differentiating omental human preadipocytes: from de-activation to generation of cortisol. *Endocr. Res.*2002;28:449-61.
35. Tomlinson JW, Sinha B, Bujalska I, Hewison M, Stewart PM. Expression of 11beta-hydroxysteroid dehydrogenase type 1 in adipose tissue is not increased in human obesity. *J. Clin. Endocrinol. Metab.* 2002; 87:5630-35.
36. Tomlinson JW, Moore J, Cooper MS, Bujalska I., Shahmanesh M, Burt C, Strain A, Hewison M, Stewart PM. Regulation of expression of 11-hydroxysteroid dehydrogenase type 1 in adipose tissue: tissue-specific induction by cytokines *Endocrinology.* 2001; 142:1982-89.
37. Franks PW, Knowler WC, Nair S, Koska J, Lee YH, Lindsay RS, et al. Interaction between an 11betaHSD1 gene variant and birth era modifies the risk of hypertension in Pima Indians. *Hypertension.*2004; 44: 681–88.
38. Caramelli E, P Strippoli, T DI Giacomi, et al. Lack of mutations of type 1 11[beta]-hydroxysteroid dehydrogenase gene in patients with abdominal obesity. *Endocr. Res.* 2001;27: 47–61.
39. Draper N, SM Echwald, GG Lavery, et al. Association studies between microsatellite markers within the gene encoding human 11betahydroxysteroid dehydrogenase type 1 and body mass index, waist to hip ratio, and glucocorticoid metabolism. *J. Clin. Endocrinol. Metab.*2002; 87: 4984–90.
40. Gelernter-Yaniv L, Feng N, Sebring NG, Hochberg Z, Yanovski JA. Associations between polymorphism in the 11 beta hydroxysteroid dehydrogenase type I gene and body composition. *Int. J. Obes. Relat. Metab. Disord* 2003; 27: 983–86.
41. Schadt EE, Lamb J, Yang X, Zhu J, Edwards S, Guhathakurta D, et al. An integrative genomics approach to infer causal associations between gene expression and disease. *Nat. Genet* 2005; 37: 710–17.
42. Otieno CJ, Bastiaansen J, Ramos AM, Rothschild MF. Mapping and association studies of diabetes related genes in the pig. *Anim. Genet* 2005; 36:36–42.
43. Reus VI, OM Wolkowitz. Antigluco-corticoid drugs in the treatment of depression. *Expert Opin. Investig. Drugs.*2001;10:1789–96.
44. Reus VI, Wolkowitz OM. Antigluco-corticoid treatment of depression: double-blind ketoconazole. *Biol. Psychol.*1999; 45: 1070–74.
45. Gaillard RC, Poffet D, Riondel AM, Saurat JH. RU486 inhibits peripheral effects of glucocorticoids in humans. *J. Clin. Endocrinol. Metab* 1985; 61:1009–11.
46. Jacobson P B, von Geldern T W, Ohman L, Osterland M, Wang J, Zinker B, et al. Hepatic glucocorticoid receptor antagonism is sufficient to reduce elevated hepatic glucose output and improve glucose control in animal models of type 2 diabetes. *J. Pharmacol. Exp. Ther* 2005; 314: 191–200.
47. Stewart P M, A M Wallace, S M Atherden, et al. Mineralocorticoid activity of carbenoxolone: contrasting effects of carbenoxolone and liquorice on 11 beta hydroxysteroid dehydrogenase activity in man. *Clin. Sci* 1990;78: 49– 54.
48. Andrew R, K Smith, GC Jones, B R Walker. Distinguishing the activities of 11 beta hydroxysteroid dehydrogenases *in vivo* using

- isotopically labeled cortisol. *J. Clin. Endocrinol. Metab* 2002; 87: 277–285.
49. Andrews R C, O Rooyackers, B R Walker. Effects of the 11 beta hydroxysteroid dehydrogenase inhibitor carbenoxolone on insulin sensitivity in men with type 2 diabetes. *J. Clin. Endocrinol. Metab* 2003; 88: 285–91.
50. Livingstone DE, Walker BR. Is 11 beta hydroxysteroid dehydrogenase type 1 a therapeutic target? Effects of carbenoxolone in lean and obese Zucker rats. *J. Pharmacol. Exp. Ther* 2003; 305: 167–72.
51. Li RS, Nakagawa Y, Nakanishi T, Fujisawa Y, Ohzeki T. Different responsiveness in body weight and hepatic 11beta-hydroxysteroid dehydrogenase (11beta-HSD) type 1 mRNA to 11beta-HSD inhibition by glycyrrhetic acid treatment in obese and lean Zucker rats. *Metabolism* 2004; 53: 600–606.
52. Barf T, Vallgarda J, Emond R, Häggström C, Kurz G, Nygren A, *et al.* Arylsulfonamidothiazoles as a new class of potential antidiabetic drugs. Discovery of potent and selective inhibitors of the 11 beta hydroxysteroid dehydrogenase type 1. *J. Med. Chem* 2002; 45: 3813–15.
53. Xiang J, Ipek M, Suri V, Masefski W, Pan N, Ge Y, *et al.* Synthesis and biological evaluation of sulfonamidooxazoles and keto sulfones: selective inhibitors of 11 β -hydroxysteroid dehydrogenase type 1. *Bioorg. Med. Chem. Lett* 2005; 15: 2865–69.
54. Hult M, H Jorvall, U C T Ppermann. Selective inhibition of human type 1 11 β -hydroxysteroid dehydrogenase by synthetic steroids and xenobiotics. *Fed. Eur. Biochem. Stud.* 1998; 441: 25–28.
55. Alberts P, L Engblom, N Edling, *et al.* Selective inhibition of 11 β hydroxysteroid dehydrogenase type 1 decreases blood glucose concentrations in hyperglycaemic mice. *Diabetologia.* 2002; 45: 1528–32.
56. Alberts P, Nilsson C, Selen G, Engblom LO, Edling NH, Norling S, *et al.* Selective inhibition of 11 beta hydroxysteroid dehydrogenase type 1 improves hepatic insulin sensitivity in hyperglycaemic mice strains. *Endocrinology.* 2003; 144: 4755–62.
57. Hermanowski-Vosatka A, Balkovec JM, Cheng K, Chen HY, Hernandez M, Koo GC, *et al.* 11 beta HSD1 inhibition ameliorates metabolic syndrome and prevents progression of atherosclerosis in mice. *J. Exp. Med* 2005; 202: 517–27.
58. Walker BR, Yau JL, Brett LP, Seckl JR, Monder C, Williams BC, *et al.* 11 β -Hydroxysteroid dehydrogenase in vascular smooth muscle and heart: implications for cardiovascular responses to glucocorticoids. *Endocrinology* 1991; 129: 3305–12.
59. Hadoke PWF, C Christy, YV Kotelevtsev, B. C. Williams, C. J. Kenyon, J. R. Seckl, *et al.* Endothelial cell dysfunction in mice after transgenic knockout of type 2, but not type 1, 11 β -hydroxysteroid dehydrogenase. *Circulation* 2001; 104: 2832–37.
60. Small G R, Hadoke P W, Sharif I, Dover A R, Armour D, Kenyon C J, *et al.* Preventing regeneration of glucocorticoids by 11 beta hydroxysteroid dehydrogenase type 1 enhances angiogenesis. *Proc. Natl. Acad. Sci. USA* .2005; 102: 12165–70
61. Wang MS, Shi H, Wang KS, Reidenberg MM. Inhibition of 11 β -hydroxysteroid dehydrogenase in guinea pig kidney by three bioflavonoids and their interactions with gossypol. *Acta Pharmacol. Sin.* 2002; 23: 92–96.
62. Thieringer R, Le Grand CB, Carbin L, Cai TQ, Wong B, Wright SD, *et al.* 11 β -Hydroxysteroid dehydrogenase type 1 is induced in human monocytes upon differentiation to macrophages. *J. Immunol.* 2001; 167: 30–35.
63. Walker BR, Yau JL, Brett LP, Seckl JR, Monder C, Williams BC, *et al.* 11 beta Hydroxysteroid dehydrogenase in vascular smooth muscle and heart: implications for cardiovascular responses to glucocorticoids. *Endocrinology* .1991; 129: 3305–12.
64. Hadoke PWF, C Christy, Y V Kotelevtsev, *et al.* Endothelial cell dysfunction in mice after transgenic knockout of type 2, but not type 1, 11 beta hydroxysteroid dehydrogenase. *Circulation.* 2001; 104: 2832–37.
65. Small GR, Hadoke PW, Sharif I, Dover AR, Armour D, Kenyon CJ, *et al.* Preventing regeneration of glucocorticoids by 11 beta hydroxysteroid dehydrogenase type 1 enhances angiogenesis. *Proc. Natl. Acad. Sci. USA.* 2005; 102: 12165–70.
66. Harris HJ, YV Kotelevtsev, JJ Mullins, *et al.* 11 beta Hydroxysteroid dehydrogenase type 1 null mice have altered hypothalamic-pituitary-adrenal axis activity: a novel control of glucocorticoid feedback. *Endocrinology.* 2001; 142: 114–20.
67. Yau JL, Noble J, Kenyon CJ, Hibberd C, Kotelevtsev Y, Mullins JJ, *et al.* Lack of tissue glucocorticoid reactivation in 11 beta hydroxysteroid dehydrogenase type 1 knockout mice ameliorates age-related learning impairments. *Proc. Natl. Acad. Sci. USA* .2001; 98: 4716–21.

68. Sandeep TC, Yau JL, MacLulich AM, Noble J, Deary IJ, Walker BR, *et al.* 11 beta hydroxysteroid dehydrogenase inhibition improves cognitive function in healthy elderly men and type 2 diabetics. *Proc. Natl. Acad. Sci. USA.*2004;101: 6734–39.
69. Stulnig TM, Oppermann U, Steffensen KR, Schuster GU, Gustafsson JA. Liver X receptors downregulate 11beta-hydroxysteroid dehydrogenase type 1 expression and activity. *Diabetes* 2002; 51:2426-33.
70. Cao G, Liang Y, Broderick CL, Oldham BA, Beyer TP, Schmidt RJ, *et al* Antidiabetic action of a liver x receptor agonist mediated by inhibition of hepatic gluconeogenesis. *J. Biol.Chem* 2003; 278:131-36.
71. Laffitte BA, Chao LC, Li J, Walczak R, Hummasti S, Joseph SB, *et al.* Activation of liver X receptor improves glucose tolerance through coordinate regulation of glucose metabolism in liver and adipose tissue. *Proc. Natl. Acad.Sci. USA* 2003; 100: 5419-24.
72. Kurukulasuriya R, Link JT, Madar DJ, Pei Z, Richards SJ, Rohde J, Souers AJ, Szczepankiewicz BG. Potential drug targets and progress towards pharmacological inhibition of hepatic glucose production. *Curr. Med.Chem* 2003; 10:123-53.
73. Link JT. Pharmacological regulation of hepatic glucose production. *Curr. Opin. Investig. Drugs* 2003; 4:421-29.
74. Jacobson PB, von Geldern TW, Ohman L, Osterland M, Wang J, Zinker B, *et al.* Hepatic glucocorticoid receptor antagonism is sufficient to reduce elevated hepatic glucose output and improve glucose control in animal models of type 2 diabetes. *J. Pharmacol. Exp. Ther* 2005; 314:191-200.
75. Valsamakis G, Anwar A, Tomlinson JW, Shackleton CH, McTernan PG, Chetty R, *et al.* 11beta-hydroxysteroid dehydrogenase type 1 activity in lean and obese males with type 2 diabetes mellitus. *J. Clin. Endocrinol. Metab* 2004; 89: 4755-61.
76. Lakshmi V, Sakai RR, McEwen BS, Monder C. Regional Distribution of 11 β -Hydroxysteroid Dehydrogenase in Rat Brain *Endocrinology* 1991; 128: 1741-48.
77. Basu R, Singh R J, Basu A, Chittilapilly E G, Johnson C M, Toffolo G, *et al.* Splanchnic cortisol production occurs in humans – evidence for conversion of cortisone to cortisol via the 11 beta hydroxysteroid dehydrogenase type 1 pathway. *Diabetes* 2004;53: 2051–59.
78. Liu Y, Nakagawa Y, Wang Y, Li R, Li X, *et al.* Leptin activation of corticosterone production in hepatocytes may contribute to the reversal of obesity and hyperglycemia in leptin-deficient ob/ob mice. *Diabetes* 2003; 52: 1409–16.
79. Tomlinson JW, Moore JS, Clark PM, Holder G, Shakespeare L, Stewart PM. Weight loss increases 11beta-hydroxysteroid dehydrogenase type 1 expression in human adipose tissue. *J. Clin. Endocrinol. Metab* 2004; 89: 2711–16.
80. Engeli S, Böhnke J, Feldpausch M, Gorzelniak K, Heintze U, Janke J, *et al.* Regulation of 11 beta hydroxysteroid dehydrogenase genes in human adipose tissue: influence of central obesity and weight loss. *Obes. Res.*2004; 12: 9–17.
81. De Kloetm E.R. Stress in brain. *Eur. J. Pharmacol* 2000; 405: 187-198.
82. De Kloet E.R, Hormones and the stressed brain. *Ann. N.Y. Acad. Sci.*2004; 1018: 1-15.
83. De Kloet, ER Vreugdenhil, E Oitzl, M.S, Joels M. Brain corticosteroid receptor balance in health and disease. *Endocr. Rev* 1998; 19: 269-301.
84. Moisan M P, Seckl J R. Edwards CR. 11 β -Hydroxysteroid Dehydrogenase Bioactivity And Messenger RNA Expression In Rat Forebrain: Localization In Hypothalamus, Hippocampus, And Cortex. *Endocrinology* 1990; 127: 1450-1455.
85. Low S C, Moisan M P, Noble J M, Edwards C R, Seckl J R. Glucocorticoids and chronic stress upregulate 11 β -hydroxysteroid dehydrogenase activity and gene expression in the hippocampus. *J. Neuroendocrinol.*1994; 6: 285-90.
86. Rajan V, Edwards CR, Seckl JR. 11 beta-Hydroxysteroid dehydrogenase in cultured hippocampal cells reactivates inert 11-dehydrocorticosterone, potentiating neurotoxicity. *J. Neurosci* 1996;16: 65-70.
87. Yau JL, Noble J, Kenyon CJ, Hibberd C, Kotelevtsev Y, Mullins JJ, Seckl JR. Lack of tissue glucocorticoid reactivation in 11beta -hydroxysteroid dehydrogenase type 1 knockout mice ameliorates age-related learning impairments. *Proc. Natl. Acad. Sick. USA* 2001;98: 4716-21.
88. Sandeep T C, Yau J L, MacLulich A M, Noble J, Deary I J, Walker B R, Seckl J R. 11 β - hydroxysteroid dehydrogenase inhibition improves cognitive function in healthy elderly men and type 2 diabetics. *Proc. Natl. Acad. Sci. USA* 2004; 101: 6734-39.
89. De Quervain D J, Poirier R, Wollmer M A, Grimaldi L M, Tsolaki M, Streffer J R, *et al.* Glucocorticoid-related genetic susceptibility

- for Alzheimer's disease. *Hum. Mol. Genet* 2004; 13: 47-52.
90. Rauz S, Walker EA, Shackleton CH, Hewison M, Murray PI, Stewart PM. Invest. Expression and putative role of 11 α -hydroxysteroid dehydrogenase isozymes within the human eye. *Ophthalmol.Vis. Sci* 2001; 42:2037-42.
91. Rauz S, Cheung C M, Wood P J, Coca-Prados M, Walker E A, Murray P I, *et al* . *Inhibition* of 11beta-hydroxysteroid dehydrogenase type 1 lowers intraocular pressure in patients with ocular hypertension. *QJM* 2003; 96:481-90.
92. Hench PS. Nobel Lecture, http://nobelprize.org/medicine/? wp_ml=0.
93. Barnes P J, Adcock I M. Steroid resistance in asthma *QJM* 1995; 88: 455-68.
94. Wilckens T. Glucocorticoids and immune function: physiological relevance and pathogenic potential of hormonal dysfunction *Trends Pharmacol. Sci* 1995; 16:193-97.
95. Wilckens T, De Rijk R. Glucocorticoids and immune function: unknown dimensions and new frontiers. *Immunol. Today* 1997; 18: 418-24.
96. Munck A, Naray-Fejes-Toth A. The ups and downs of glucocorticoid physiology. Permissive and suppressive effects revisited. *Mol. Cell Endocrinol.* 1992;90: C1-4.
97. Zhou Z, Shackleton C H, Pahwa S, White P C, Speiser P W. Prominent sex steroid metabolism in human lymphocytes. *Mol. Cell Endocrinol.* 1998; 138: 61-69.
98. Hennebold J D, Ryu S Y, Mu H H, Galbraith A, Daynes R A. 11 β -hydroxysteroid dehydrogenase modulation of glucocorticoid activities in lymphoid organs. *Am. J. Physiol* 1996; 270: R1296-1306.
99. Freeman L, Hewison M, Hughes SV, Evans KN, Hardie D, Mean TK. Expression of 11beta-hydroxysteroid dehydrogenase type 1 permits regulation of glucocorticoid bioavailability by human dendritic cells. *Blood* 2005; 106: 2042-49.
100. Teelucksingh S, Mackie A D, Burt D, McIntyre M A, Brett L, Edwards CR. Potentiation of hydrocortisone activity in skin by glycyrrhetic acid. *Lancet* 1990; 335:1060-63.
101. Hennebold J D, Mu H H, Poynter M E, Chen X P, Daynes RA. Active catabolism of glucocorticoids by 11 beta-hydroxysteroid dehydrogenase in vivo is a necessary requirement for natural resistance to infection with *Listeria monocytogenes*. *Int. Immunol* 1997; 9: 105-115.
102. Manelli F, Giustina A. Glucocorticoid-induced osteoporosis. *Metab* 2000; 11: 79-85.
103. Doga M, Bonadonna S, Burattin A, Carpinteri R, Manelli F, Giustina A. Bisphosphonates in the Treatment of Glucocorticoid-Induced Osteoporosis. *Front. Horm. Res* .2002; 30:150-164.
104. Cooper M S, Rabbitt E H, Goddard P E, Bartlett W A, Hewison M, Stewart P M. Osteoblastic 11beta-hydroxysteroid dehydrogenase type 1 activity increases with age and glucocorticoid exposure. *J. Bone Miner. Res* 2002; 17: 979-86.
105. Cooper MS, Walker EA, Bland R, Fraser WD, Hewison M, Stewart PM. Expression and functional consequences of 11beta-hydroxysteroid dehydrogenase activity in human bone. *Bone*.2000; 27:375-81.
106. Cooper MS, Blumsohn A, Goddard PE, Bartlett WA, Shackleton CH, Eastell R, *et al*. 11 β hydroxysteroid dehydrogenase type 1 activity predicts the effects of glucocorticoids on bone. *J. Clin. Endocrinol. Metab* 2003; 88:3874-77.
107. Justesen J, Mosekilde L, Holmes M, Stenderup K, Gasser J, Mullins J J, *et al*. Mice deficient in 11beta-hydroxysteroid dehydrogenase type 1 lack bone marrow adipocytes, but maintain normal bone formation. *Endocrinology* 2004.145:1916-25.